



Seasonal features of geomagnetic activity: a study on the solar activity dependence

Adriane Marques de Souza Franco¹, Rajkumar Hajra², Ezequiel Echer¹, and Mauricio José Alves Bolzan³

¹Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, Brazil

²Indian Institute of Technology Indore, Simrol, Indore 453552, India

³Space Physics and Astronomy Laboratory, Federal University of Jatai, Jatai, Brazil

Correspondence: Adriane Marques de Souza Franco (adrianemarquesds@gmail.com) and Rajkumar Hajra (rajkumarhajra@yahoo.co.in)

Received: 24 May 2021 – Discussion started: 24 June 2021

Revised: 16 September 2021 – Accepted: 17 September 2021 – Published: 18 October 2021

Abstract. Seasonal features of geomagnetic activity and their solar-wind–interplanetary drivers are studied using more than five solar cycles of geomagnetic activity and solar wind observations. This study involves a total of 1296 geomagnetic storms of varying intensity identified using the Dst index from January 1963 to December 2019, a total of 75 863 substorms identified from the SuperMAG AL/SML index from January 1976 to December 2019 and a total of 145 high-intensity long-duration continuous auroral electrojet (AE) activity (HILDCAA) events identified using the AE index from January 1975 to December 2017. The occurrence rates of the substorms and geomagnetic storms, including moderate ($-50 \text{ nT} \geq \text{Dst} > -100 \text{ nT}$) and intense ($-100 \text{ nT} \geq \text{Dst} > -250 \text{ nT}$) storms, exhibit a significant semi-annual variation (periodicity ~ 6 months), while the super storms ($\text{Dst} \leq -250 \text{ nT}$) and HILDCAAs do not exhibit any clear seasonal feature. The geomagnetic activity indices Dst and ap exhibit a semi-annual variation, while AE exhibits an annual variation (periodicity ~ 1 year). The annual and semi-annual variations are attributed to the annual variation of the solar wind speed V_{sw} and the semi-annual variation of the coupling function VB_s (where $V = V_{\text{sw}}$, and B_s is the southward component of the interplanetary magnetic field), respectively. We present a detailed analysis of the annual and semi-annual variations and their dependencies on the solar activity cycles separated as the odd, even, weak and strong solar cycles.

1 Introduction

Solar-wind–magnetosphere energy coupling causes disturbances in the magnetosphere of the Earth (e.g., Dungey, 1961; Axford and Hines, 1961; Tsurutani et al., 1992; Gonzalez et al., 1994; Tsurutani et al., 2020). Depending on the strength, duration and efficiency of the coupling, resultant geomagnetic disturbances (von Humboldt, 1808) can be classified as magnetic storms, substorms and high-intensity long-duration continuous auroral electrojet (AE) activities (HILDCAAs) (see Gonzalez et al., 1994; Hajra et al., 2020; Hajra, 2021a). In general, magnetic storms represent global-scale disturbances caused by enhancements in (westward) ring current flowing at $\sim 2\text{--}7$ Earth radii (R_{Earth}) in the magnetic equatorial plane of the Earth (Gonzalez et al., 1994; Lakhina and Tsurutani, 2018, and references therein). Storm duration spans a few hours to several days. In fact, while the storm main phase lasts typically for $\sim 10\text{--}15$ h, the recovery phase can continue much longer, from hours to several days (Gonzalez et al., 1994). Substorms (Akasofu, 1964) are shorter-scale, a few minutes to a few hours, disturbances in the night-side magnetosphere (magnetotail) resulting in precipitations of $\sim 10\text{--}100$ keV electrons and protons in the auroral ionosphere (e.g., Meng et al., 1979; Thorne et al., 2010; Tsurutani et al., 2019, and references therein). Intense auroral substorms continuing for a few days without occurrence of any major magnetic storms have been called HILDCAAs (Tsurutani and Gonzalez, 1987; Hajra et al., 2013) to distinguish them from nominal substorms and major magnetic storms (Tsurutani et al., 2004; Guarnieri, 2006).

It is important to note that from the physical point of view, substorms and HILDCAAs are two different types of geomagnetic activity. While substorms may occur during HILDCAAs, they represent different magnetosphere/ionosphere processes (Tsurutani et al., 2004; Guarnieri, 2005, 2006). For example, HILDCAAs are associated with Alfvén wave trains carried by solar wind high-speed ($\sim 550\text{--}850\text{ km s}^{-1}$) streams (HSSs) emanated from solar coronal holes (Tsurutani and Gonzalez, 1987; Hajra et al., 2013). The intermittent magnetic reconnection between the Alfvén wave southward component and geomagnetic field results in intermittent increases in auroral activity during HILDCAAs. Substorms, on the other hand, are associated with solar wind energy loading in the magnetotail caused by magnetic reconnection (Tsurutani and Meng, 1972), and subsequent explosive release of the energy in the form of energetic particles and strong plasma flows (e.g., Akasofu, 1964, 2017; Rostoker, 2002; Nykyri et al., 2019, and references therein). These are not essentially associated with HSSs. Thus, for good reason, the term “substorm” was avoided in the definition of HILDCAAs by Tsurutani and Gonzalez (1987). Later, Hajra et al. (2014b, 2015a, b) have shown that HILDCAAs take an important role in the acceleration of relativistic ($\sim \text{MeV}$) electrons in the outer radiation belt of the Earth. This feature further distinguishes the HILDCAAs from nominal substorms.

Geomagnetic activity, in general, is known to be highly variable, modulated by several solar–terrestrial features. The solar/interplanetary sources of the variability include the $\sim 27\text{ d}$ solar rotation (Bartels, 1932, 1934; Newton and Nunn, 1951), the $\sim 11\text{-year}$ solar activity cycle (Schwabe, 1844), the electromagnetic and corpuscular radiations from the Sun, several plasma emission phenomena, heliospheric current region, etc. On the other hand, the Earth’s translational movement (solstices), the inter-hemispheric symmetry (equinoxes), and the observational frame of reference or the coordinate system (Russell, 1971) can also largely impact the geomagnetic activity variation.

One of the earliest reported features of the geomagnetic activity is the semi-annual variation, that is, more frequent occurrences and higher strength during equinoxes and lesser occurrences and weaker strength during solstices (e.g., Broun, 1848; Sabine, 1852). The semi-annual variation is reported in the occurrence rates and intensities of the magnetic storms (e.g., Cliver et al., 2000, 2004; Le Mouél et al., 2004; Cnossen and Richmond, 2012; Danilov et al., 2013; McPherron and Chu, 2018; Lockwood et al., 2020) and in the Earth’s radiation belt electron variations (e.g., Baker et al., 1999; Li et al., 2001; Kanekal et al., 2010; Katsavrias et al., 2021). This is generally explained in the context of the Earth’s position in the heliosphere (known as the “axial effect”; Cortie, 1912), relative angle of solar wind incidence with respect to Earth’s rotation axis (the “equinoctial effect”; Boller and Stolov, 1970) and geometrical controls of interplanetary magnetic fields (the “Russell–McPherron effect”; Russell and McPherron, 1973). See Lockwood et al. (2020) for

an excellent discussion of the mechanisms. While both the equinoctial and the Russell–McPherron effects are shown to be responsible for the semi-annual variation in the geomagnetic indices (e.g., Cliver et al., 2000; O’Brien and McPherron, 2002), the semi-annual variation in the relativistic electron fluxes of the outer belt is mainly attributed to the Russell–McPherron effect (e.g., Kanekal et al., 2010; Katsavrias et al., 2021).

However, the semi-annual variation in general was questioned by the work of Mursula et al. (2011) reporting solstice maxima in substorm frequency and duration, as well as substorm amplitude and global geomagnetic activity peaks alternating between spring and fall in ~ 11 years. While solstice maxima were attributed to auroral ionospheric conductivity changes (Wang and Lühr, 2007; Tanskanen et al., 2011), the alternating equinoctial maxima were associated with asymmetric solar wind distribution in solar hemispheres (Mursula and Zieger, 2001; Mursula et al., 2002). In addition, several recent studies have reported a lack of any seasonal dependence for substorms (Hajra et al., 2016), HILDCAAs (Hajra et al., 2013, 2014a) or in the radiation belts (Hajra, 2021b).

In the present work, for the first time, we will explore a long-term database of substorms, HILDCAAs, and magnetic storms of varying intensity along with different geomagnetic indices to study the seasonal features of geomagnetic disturbances. The main aim is to identify and characterize the seasonal features of geomagnetic disturbances of different types and intensities. In addition, we will study their solar activity dependencies, if any.

2 Database and methods

Details of the geomagnetic events studied in this work are summarized in Table 1. Auroral substorms are identified by intensification in the auroral ionospheric (westward) electrojet currents. In the present work, we will use the substorm list available at the SuperMAG website (<https://supermag.jhuapl.edu/>, last access: 23 May 2021; Newell and Gjerloev, 2011; Gjerloev, 2012). The substorm expansion phase onsets were identified from the SML index which is the SuperMAG equivalent of the westward auroral electrojet index AL (see the cited references for details). The present work involves a total of 75 863 substorms identified from January 1976 to December 2019 (Table 1).

We will use the geomagnetic storm and HILDCAA database prepared by Hajra et al. (2021) for the present work. It is an updated version of the lists presented in Echer et al. (2011), Hajra et al. (2013) and Rawat et al. (2018). Geomagnetic storm onset, main phase, peak strength, recovery phase and storm end are determined by the variations of the Dst index (Sugiura, 1964). Based on the Gonzalez et al. (1994) definition, intervals with the Dst minimum $\leq -50\text{ nT}$ are identified as magnetic storms. From January 1963 to December 2019, 1296 magnetic storms were identified (Table 1).

Table 1. Details of the geomagnetic activity events under present study.

Events	Number of events	Periods of observation	Geomagnetic indices	Sources of events
Substorms	75 863	January 1976–December 2019	SML	https://supermag.jhuapl.edu/ (last access: 23 May 2021)
HILDCAAs	145	January 1975–December 2017	AE, Dst	Hajra et al. (2021)
Geomagnetic storms	1296	January 1963–December 2019	Dst	Hajra et al. (2021)

Geomagnetic storms with the Dst minimum values between -50 nT and -100 nT are classified as the “moderate storms”, between -100 nT and -250 nT as the “intense storms”, and those with the Dst minima lower than -250 nT as the “super storms”. Among all storms studied here, 75 % are moderate, 23 % are intense and only 2 % are super storms.

The HILDCAA events are identified based on four criteria suggested by Tsurutani and Gonzalez (1987). The criteria are (1) the AE index should reach an intensity equal to or greater than 1000 nT at some point during the event (the high-intensity criterion), (2) the event must last at least 2 d (the long-duration criterion), (3) the AE index should not fall below 200 nT for more than 2 h at a time (the continuity criterion), and (4) the auroral activity must occur outside the main phase of a geomagnetic storm or during a non-storm condition ($Dst > -50$ nT). Present work involves a total of 145 HILDCAA events identified during January 1975 through to December 2017 (Table 1).

The geomagnetic indices, namely, the ring current index Dst, the global-scale geomagnetic activity index ap and the auroral ionospheric current-related index AE, are used to provide a quantitative measure of the activity level of the terrestrial magnetosphere (Rostoker, 1972). In addition, solar wind parameters are used to study the energy dissipation in the magnetosphere. The D_{500} parameter is defined as the percentage of days with the peak solar wind speed V_{sw} equal or higher than 500 km s^{-1} in each month of a year. We estimated the solar wind electric field VB_s , which is an important solar-wind–magnetosphere coupling function (Burton et al., 1975; Tsurutani et al., 1992; Finch et al., 2008). As VB_s involves both the solar wind velocity V_{sw} (for V) and the southward component of the interplanetary magnetic field (IMF) B_s , the latter being important for magnetic reconnection, VB_s is also called the reconnection electric field. The Akasofu ϵ coupling function (Perreault and Akasofu, 1978), expressed as $V_{sw} B_0^2 \sin^4(\theta/2) R_{CF}^2$, was also estimated in this work as a proxy for the magnetospheric energy input rate. Here B_0 represents the magnitude of the IMF, θ is the IMF orientation clock angle and R_{CF} is the Chapman–Ferraro magnetopause distance (Chapman and Ferraro, 1931).

The 10.7 cm solar flux ($F_{10.7}$) is shown to be a good indicator of the solar activity (e.g., Tapping, 1987). Thus, the ~ 11 -year solar cycles (Schwabe, 1844) are studied using the monthly mean $F_{10.7}$ solar flux variation. The starting, peak and end dates along with the peak $F_{10.7}$ flux of each solar cycle (SC) are listed in Table 2. The $F_{10.7}$ fluxes are given in the solar flux unit (sfu), where $1 \text{ sfu} = 10^{-22} \text{ W m}^{-2} \text{ Hz}^{-1}$.

Table 2. Details of the solar cycles under present study.

SC no.	SC start date	SC peak date	SC peak $F_{10.7}$	SC end date
SC20	Oct 1964	Nov 1968	156	Feb 1976
SC21	Mar 1976	Dec 1979	203	Aug 1986
SC22	Sep 1986	Nov 1989	213	Jul 1996
SC23	Aug 1996	Nov 2001	181	Nov 2008
SC24	Dec 2008	Apr 2014	146	Dec 2019

Based on the $F_{10.7}$ peaks, cycles SC20 and SC24 can be classified as the “weak cycles” (average $F_{10.7}$ peak ~ 151 sfu) and SC19, SC21, SC22 and SC23 as the “strong cycles” (average $F_{10.7}$ peak ~ 207 sfu). It can be mentioned that SC24 is the weakest cycle in the space exploration era (after 1957). A detailed study on the solar and geomagnetic characteristics of this cycle is presented in Hajra (2021c). The solar cycles are also grouped into the “even” (SC20, SC22, SC24) and the “odd” (SC19, SC21, SC23) cycles in this work. Previous studies have reported significant differences between the even and odd cycle amplitudes (e.g., Waldmeier, 1934; Gnevyshev and Ohl, 1948; Wilson, 1988; Durney, 2000), and in their geomagnetic responses (e.g., Hajra et al., 2021; Owens et al., 2021).

We will apply the Lomb–Scargle periodogram analysis (Lomb, 1976; Scargle, 1982) to identify the significant periodicities in the geomagnetic event occurrences, the geomagnetic indices and the solar-wind–magnetosphere (coupling) parameters. It is a useful tool for detecting and characterizing periodic signals for unequally spaced data.

The geomagnetic indices are collected from the World Data Center for Geomagnetism, Kyoto, Japan (<http://wdc.kugi.kyoto-u.ac.jp/>, last access: 23 May 2021). The monthly means of the solar wind/interplanetary data near the Earth’s bow shock nose were obtained from NASA’s OMNI database (<http://omniweb.gsfc.nasa.gov/>, last access: 23 May 2021). The IMF vector components are in Geocentric Solar Magnetospheric (GSM) coordinates, where the x axis is directed towards the Sun and the y axis is in the $\mathbf{\Omega} \times \hat{\mathbf{x}}/|\mathbf{\Omega} \times \hat{\mathbf{x}}|$ direction, where $\mathbf{\Omega}$ is aligned with the magnetic south pole axis of the Earth, and $\hat{\mathbf{x}}$ is the unit vector along the x axis. The z axis completes a right-hand system. The $F_{10.7}$ solar fluxes are obtained from the Laboratory for Atmospheric and Space Physics (LASP) Interactive Solar Irradiance Data Center (<https://lasp.colorado.edu/lisird/>, last access: 23 May 2021).

3 Results

3.1 Seasonal features

Figure 1 shows the variations of the monthly mean solar $F_{10.7}$ flux (Fig. 1a); the monthly numbers of HILDCAAs and substorms (Fig. 1b); magnetic storms of varying intensity (Fig. 1c); the monthly mean geomagnetic Dst (Fig. 1d), ap (Fig. 1e), and AE (Fig. 1f) indices; the IMF magnitude B_0 (Fig. 1g); the solar wind plasma speed V_{sw} (Fig. 1h); the percentage occurrences of $V_{sw} \geq 500 \text{ km s}^{-1}$ (D_{500} , Fig. 1i); and the energy coupling functions VB_s (Fig. 1j) and ϵ (Fig. 1k) for the period from 1963 to 2019. While most of the data span for more than five solar cycles, from the beginning of SC20 to the end of SC24, substorm and HILDCAA data are only available from SC21 onward. The $F_{10.7}$ solar flux variation shows a clear ~ 11 -year solar activity cycle, with the minimum flux during the solar minimum, followed by flux increases during the ascending phase leading to the peak flux during the solar maximum, and flux decreases during the descending phase of the solar cycle (Fig. 1a). In general, the substorm, HILDCAA, and geomagnetic storm numbers; the geomagnetic indices; and the solar wind parameter values exhibit an overall ~ 11 -year periodicity. Embedded in the large-scale ~ 11 -year variations, there are several short-term fluctuations in the data; some of the latter may be associated with the annual or semi-annual variations, which will be explored in detail in the following sections.

Monthly superposed variations

Figure 2 shows the monthly superposed values of all the parameters shown in Fig. 1. Figure 2a–f shows the numbers of geomagnetic events in each month divided by the number of years of observations (in the unit of number per year). Figure 2g–l shows the monthly means of the geomagnetic and solar wind/interplanetary parameters for the entire interval of study.

The substorm occurrence rate (Fig. 2a) clearly exhibits two peaks during the months of March and October and a summer solstice minimum (during the month of June). HILDCAAs (Fig. 2b) do not exhibit any clear seasonal feature, except a significant minimum in November. Geomagnetic storms, from moderate to intense (Fig. 2d–e), exhibit a clear semi-annual variation. The spring equinoctial peak is recorded during March for the moderate storms and during April for the intense storms, while the fall peak is recorded during October for both of them. The super storms (Fig. 2f), with a very low occurrence rate, do not have any clear seasonal feature. As majority of the storms are of moderate intensity; storms of all intensity together (Fig. 2c) exhibit a prominent semi-annual variation with two peaks during March and October.

The monthly mean intensities of the Dst (Fig. 2g) and ap (Fig. 2h) indices show a semi-annual variation. Both of them exhibit the spring peaks during March. While Dst has




Figure 1. From top to bottom, the panels show (a) the monthly mean solar $F_{10.7}$ flux; (b) monthly numbers of substorms (black, legend on the left) and HILDCAAs (red, legend on the right); (c) geomagnetic storms of varying intensity; monthly mean (d) Dst, (e) ap, (f) AE, (g) IMF B_0 , and (h) V_{sw} ; (i) percentage of days with daily peak $V_{sw} \geq 500 \text{ km s}^{-1}$ (D_{500}); and monthly mean (j) VB_s and (k) Akasofu ϵ parameter, during 1963 to 2020. Solar cycles from SC20 to SC24 are marked on the top panel.

a fall minimum during October, ap exhibits a peak during September. On the other hand, the monthly mean AE index (Fig. 2i) increases gradually from January; attains a peak around April; decreases with a much slower rate till September, after which the decrease rate is faster, and finally attains a minimum during December. Thus, the AE index shows an annual variation, different from the Dst and ap indices. This result is consistent with Katsavrias et al. (2016) who also reported an annual component in AE, and lack of any semi-annual component. As the AE index is based on geomagnetic observations made in the northern hemisphere, the asymmetric pole exposition to the solar radiation during the Earth's translational motion could contribute to this annual variation. The latter may modulate the AE current through the modulation of the ionospheric conductivity, owing to the solar extreme ultraviolet (EUV) ionization.




Figure 2. Monthly superposed variations. Left panels, from top to bottom, show the total numbers divided by years of observation of (a) substorms, (b) HILDCAAs, (c) all storms (AS), (d) moderate storms (MS), (e) intense storms (IS) and (f) super storms (SS), respectively. Right panels, from top to bottom, show the monthly mean values of the (g) geomagnetic Dst, (h) ap, and (i) AE indices; (j) IMF B_0 ; (k) V_{sw} (black, legend on the left) and D_{500} (red, legend on the right); and (l) VB_s (black, legend on the left) and ϵ parameter (red, legend on the right), respectively.

It is worth mentioning that the AE index (Davis and Sugiura, 1966) includes an upper envelope (AU) and a lower envelope (AL) related to the largest (positive) and smallest (negative) magnetic deflections, respectively, among the magnetometer stations used. The AU and AL components represent the strengths of the eastward and westward AE, respectively. Lockwood et al. (2020) showed that the semi-annual variation is indeed present in the AL index. As the auroral westward current represented by AL is associated with the substorm-related energetic particle precipitation in the auroral ionosphere, the semi-annual variation in AL is consistent with the semi-annual variation exhibited by the substorms (present work). On the other hand, the eastward auroral current/AU is mainly contributed by the dayside ionospheric conductivity that exhibits a summer solstice maximum as suggested by Wang and Lühr (2007) and Tanskanen et al. (2011).

Among the solar-wind–magnetosphere coupling parameters, VB_s (Fig. 2l, legend on the left) exhibits a semi-annual variation, with larger average values during February–April months, another sharp peak during October and with a solstice minimum. For the monthly mean IMF B_0 (Fig. 2j), a clear minimum can be noted during July, and B_0 increases gradually on both sides of July. No clear seasonal features can be inferred from the variations of the monthly mean V_{sw} (Fig. 2k, legend on the left) or Akasofu ϵ parameter (Fig. 2l, legend on the right). However, D_{500} (Fig. 2k, legend on the

right) exhibits two clear peaks around March and September, with prominently lower values during solstices.

Periodogram analysis

It should be noted that the seasonal features as described above (Fig. 2) present an average scenario composed by superposition of several solar cycles. The seasonal features may vary from one solar cycle to the other. In Fig. 3 we have performed the Lomb–Scargle periodogram analysis of the above events and parameters. For this purpose, we use the monthly means of $F_{10.7}$, Dst, ap, AE, B_0 , V_{sw} , D_{500} , VB_s and ϵ , as well as the monthly numbers of substorms, HILDCAAs and magnetic storms of varying intensity. In the left panel of Fig. 3, the periodograms are based on the original data of 1-month resolution, while the right panel shows the periodograms after filtering out the dominating ~ 11 -year periodicity from the data. It can be noted that the filtering helps to better identify the shorter-scale periodicities in the time series.

As expected, the $F_{10.7}$ solar flux shows a prominent (at $> 95\%$ significance level) ~ 11 -year periodicity (Fig. 3a) and no shorter-scale variation (Fig. 3b). A dominating ~ 11 -year periodicity can also be observed in substorms; HILDCAAs (Fig. 3c); magnetic storms of varying intensity (Fig. 3e); the geomagnetic indices Dst, ap, and AE (Fig. 3g); in the solar wind/interplanetary parameters IMF B_0 , V_{sw} , and D_{500} (Fig. 3i); and the solar-wind–magnetosphere coupling functions VB_s and ϵ (Fig. 3k). However, we are interested in the




Figure 3. Lomb–Scargle periodograms. From top to bottom, the panels show the normalized power of periods for the monthly mean (a, b) solar $F_{10.7}$ flux; monthly numbers of (c, d) substorms and HILDCAAs; (e, f) all magnetic storms, moderate storms, and intense storms; monthly mean (g–h) geomagnetic indices Dst, ap, and AE; (i–j) solar wind parameters IMF B_0 , V_{sw} , and D_{500} ; and (k–l) VB_s and ϵ parameter, respectively. The left panels correspond to periodograms of the original database without any filtering, while the right panels correspond to periodograms after filtering out the 11-year periodicity from the database. Horizontal dot-dashed lines in each panel indicate $> 95\%$ significance levels of the corresponding parameters shown by different colors. Note that the x axes have different scaling for the left and right panels.

annual or shorter-scale periodicities in the events and parameters. Thus, the Lomb–Scargle periodograms are also performed after filtering out this dominating ~ 11 -year periodicity from the data. The same is shown in the right panel of Fig. 3.

Table 3 lists the significant periodicities which are less than the ~ 11 -year solar cycle period. As clear from Fig. 3 and Table 3, substorms (Fig. 3d) and moderate and intense geomagnetic storms (Fig. 3f) exhibit prominent semi-annual (~ 6 -month period) variation. However, the super storms do not exhibit any clear variation pattern (not shown). HILDCAAs (Fig. 3d), on the other hand, exhibit a ~ 4.1 -year periodicity, while no annual or lower-scale variation was recorded. However, it should be noted that very low monthly numbers of HILDCAAs and super storms during different years may introduce significant artifacts to the corresponding spectral/periodogram analysis. Thus, the results of the periodogram analysis for HILDCAAs and super storms cannot be fully trusted.

Both the ap and Dst indices exhibit a clear ~ 6 -month periodicity (Fig. 3h). However, the AE index exhibits an annual variation but no semi-annual variation.

The solar wind/interplanetary and coupling functions exhibit more complex periodicity (lower than ~ 11 years). The

IMF B_0 (Fig. 3i) and ϵ parameter (Fig. 3k) exhibit ~ 8 -year periodicity but no annual or lower-scale periodicity (Fig. 3j and l). The solar wind V_{sw} and D_{500} (Fig. 3j) exhibit several periodicities in the range of ~ 4 –8 years and a significant annual variation (periodicity ~ 1 year). The coupling function VB_s exhibits a prominent semi-annual variation (Fig. 3l). The V_{sw} periodicities detected in the present work are consistent with results reported previously (e.g., Valdés-Galicia et al., 1996; El-Borie, 2002; El-Borie et al., 2020; Hajra, 2021a; Hajra et al., 2021, and references therein). For example, El-Borie (2002) reported ~ 9.6 -year periodicity in V_{sw} arising from the coronal hole variations in the southern hemisphere of the Sun. El-Borie et al. (2020) discussed multiple V_{sw} periodicities in the 1–2-, 2–4-, 4–8- and 8–16-year bands. Recently, Hajra et al. (2021) reported significant V_{sw} periodicities of ~ 3 , ~ 4 , ~ 10 and ~ 16 years and discussed their important role in space climatology.

The results shown in Fig. 3 and Table 3 are consistent with those in Fig. 2. From the above analyses, the coupling function VB_s which exhibits a ~ 6 -month periodicity can be inferred as the driver of the semi-annual variations in substorms, moderate and intense storms, and in the geomagnetic indices Dst and ap. On the other hand, the ~ 1 -year periodicity in V_{sw}/D_{500} can be a source of the annual variation in the

Table 3. Significant (at the > 95% level) periods less than ~ 11 years obtained from the Lomb–Scargle periodogram analysis. Periods are ordered from higher power to lower.

Events/parameters	Period (year)
Geomagnetic activity	
Substorms	0.5, 4.2
HILDCAAs	4.1
All storms	0.5
Moderate storms	0.5
Intense storms	0.5
Super storms	No
Geomagnetic indices	
Dst	0.5
ap	0.5
AE	1.0
Solar wind parameters	
B_0	8.0
V_{sw}	8.3, 4.7, 1.1
D_{500}	8.3, 7.0, 5.4, 4.8, 4.3, 3.6, 1.1
VB_s	0.5
ϵ	8.1

AE index. In addition, the ~ 4.1-year periodicity in HILD-CAAs seems to be associated with the solar wind V_{sw} variation in the same range. Detailed analyses of the events and/or parameters which exhibit the annual and/or semi-annual variations are shown in Sect. 3.2. For a detailed analysis of the longer-scale variations of the geomagnetic activity, the geomagnetic indices, and the solar-wind–magnetosphere coupling, which is beyond the scope of this present work, we refer the reader to Hajra et al. (2021).

3.2 Solar activity dependence

The solar cycle variations of the seasonal features described in Sect. 3.1 are explored in Figs. 4 to 11. They show the variations of the substorms (Fig. 4); the moderate (Fig. 5) and intense (Fig. 6) magnetic storms; the geomagnetic Dst (Fig. 7), ap (Fig. 8), and AE (Fig. 9) indices; the solar wind plasma speed V_{sw} (Fig. 10); and the coupling function VB_s (Fig. 11). The format is identical for all these figures: for the geomagnetic events (the solar wind/interplanetary parameters), panel c shows the year–month contour plot of the number of the events (the mean values of the parameters) in each month of the observing years. The values of different colors are given in the legend at the bottom. Panel d shows the yearly mean $F_{10.7}$ solar flux. The solar minima are marked by the horizontal dot-dashed lines in the bottom panels c–d. Panel b shows the monthly numbers of the events per a year of observation (the monthly mean values of the parameters) during each so-

Table 4. Seasonal modulation (%) between the equinoctial maximum and the solstice minimum for the events and the parameters with the semi-annual variation during the weak and strong solar cycles, as well as the odd and even solar cycles (defined in Sect. 2).

Events/parameters	Weak solar cycle	Strong solar cycle	Odd solar cycle	Even solar cycle
Substorms	55	46	49	66
All storms	85	76	76	78
Moderate storms	92	73	68	77
Intense storms	92	100	133	105
Dst	67	85	96	79
ap	40	37	38	46
VB_s	54	57	53	40

lar cycle, while panel a shows the same during groups of the even, odd, strong, weak and all solar cycles.

Table 4 lists a “seasonal modulation” parameter defined as the difference between the equinoctial maximum and the solstice minimum expressed as the percentage of the yearly mean value for the events and parameters exhibiting the semi-annual variation. The modulation parameter can be taken as a measure of the seasonal/semi-annual variability. The larger the value of the parameter, the stronger the semi-annual variability. Large variation in the seasonal modulation can be noted from the table. For substorms, all storms, moderate storms and the ap index, seasonal modulations are larger during the weak cycles (even cycles) than the strong cycles (odd cycles). However, the modulations are larger during the strong cycles (odd cycles) than the weak cycles (even cycles) for the intense storms, the Dst index and the coupling function VB_s . The explanation is not known at present. However, it is interesting to note that the intense storms (and thus the strong Dst associated with intense VB_s) are mainly driven by the interplanetary coronal mass ejections (ICMEs). On the other hand, the moderate storms, substorms, and the ap index variations are associated with both ICMEs, and the corotating interaction regions (CIRs) between the slow streams and HSSs (e.g., Tsurutani and Gonzalez, 1987; Tsurutani et al., 1988; Gosling et al., 1990; Richardson et al., 2002; Echer et al., 2008; Hajra et al., 2013; Souza et al., 2016; Mendes et al., 2017; Marques de Souza et al., 2018; Tsurutani et al., 2019, and references therein). The strong cycles are expected to be characterized by more solar transient events like ICMEs than during the weak cycles. However, recent studies show lower numbers and reduced geoeffectiveness of both CIRs and ICMEs during the weak cycles than during the strong cycles (e.g., Scolini et al., 2018; Grandin et al., 2019; Lamy et al., 2019; Nakagawa et al., 2019; Syed Ibrahim et al., 2019; Hajra, 2021c, and references therein). This calls for a further study to explain the above results.




Figure 4. Substorms from 1976 to 2019. Panel (c) shows the year–month contour plot of the number of substorms in each month of the years 1976–2019. The values of different colors are given in the legend at the bottom. Data gaps are shown by crosses. Panel (d) shows the yearly mean $F_{10.7}$ solar flux. Panel (b) shows the monthly numbers of substorms per a year of observation during each solar cycles, and panel (a) shows the same during groups of the even, odd, strong, weak and all solar cycles. For details on the grouping of the solar cycles, see the text. The solar minima are marked by horizontal dot-dashed lines.

3.2.1 Substorms

From Fig. 4c it can be seen that in any solar cycle, the peak substorm occurrence rates are noted during the descending phase, followed by the occurrence minimum during the solar minimum to early ascending phase. From the four complete solar cycles (SC21–SC24) of the substorm observations, two prominent peaks can be noted in the years of 1994 and 2003, which are in the descending phases of SC22 and SC23, respectively.

On the seasonal basis, two peaks around the months of March and October can be observed from the year–month contour plot (Fig. 4c), which is also reflected in the monthly superposed plots (Fig. 4a–b). However, this “semi-annual” variation exhibits a large asymmetry in amplitude and duration between the spring and fall equinoxes. For example, in the year 1994, the substorm occurrence peak during February–May is significantly larger than the occurrences




Figure 5. Moderate geomagnetic storms from 1963 to 2019. The panels are in the same format as in Fig. 4.

during October. On the other hand, during 2003, while the occurrence peak is noted in November, comparable occurrences are clear almost during the entire year.

When separated on the basis of the solar cycles (Fig. 4a–b), the smallest numbers of events are observed during SC24. Interestingly, the spring occurrences are the strongest in SC22 and the fall occurrences are the strongest in SC23. Another noteworthy feature is that the occurrence rates during the even and weak solar cycles are lower than during the odd and strong cycles, respectively. However, the seasonal modulation between the equinoctial maximum and the solstice minimum is comparable between the weak ($\sim 55\%$) and strong ($\sim 46\%$) cycles (Table 4).

3.2.2 Geomagnetic storms

Variations of the moderate and intense geomagnetic storms are shown in Figs. 5 and 6, respectively. From the year–month contour plots (Figs. 5c and 6c), the moderate storms are found to peak around the descending phases, while the intense storms peak around the solar maximum. When the monthly variations of the storms are considered in each year, there is hardly any seasonal variation. However, when observations during several solar cycles are grouped together (Figs. 5a and 6a), the semi-annual variation can be noted in the moderate storms. There is not much difference in moder-




Figure 6. Intense geomagnetic storms from 1963 to 2019. The panels are in the same format as in Fig. 4.

ate and intense storm occurrence rates between the odd and even cycles. However, the occurrence rates of the storms are slightly larger in the strong cycles compared to the weak ones, while the seasonal modulation between the equinoctial maximum and the solstice minimum during the strong and weak cycles is comparable (Table 4). Another noteworthy feature is the lowest occurrence of intense storms during SC24.

3.2.3 Geomagnetic indices

Variations of the monthly mean geomagnetic indices are shown in Figs. 7 (Dst), 8 (ap) and 9 (AE). In each solar cycle, the average Dst index exhibits the strongest negative excursions at and immediately after the solar maximum (Fig. 7c–d). A clear correlation can be observed between the $F_{10.7}$ solar flux and the average Dst strength. The Dst negative excursions are stronger during the strong and odd cycles compared to the weak and even cycles, respectively (Fig. 7a). In addition, the seasonal modulation between the equinox minimum to the solstice maximum is significantly higher in the strong cycles (~ 85 %) compared to the weak cycles (~ 67 %) (Table 4). During SC24, the overall Dst strength is the weakest and there is no prominent seasonal modulation.

Variation of the monthly mean ap index (Fig. 8) is identical to the Dst index variation. However, the seasonal modulation




Figure 7. Geomagnetic Dst index variation from 1963 to 2019. Panel (c) shows the year–month contour plot of the mean Dst value in each month of the years 1963–2019. The values of different colors are given in the legend at the bottom. Data gaps are shown by crosses. Panel (d) shows the yearly mean $F_{10.7}$ solar flux. Panel (b) shows the monthly means of Dst during each solar cycles, and panel (a) shows the same during groups of the even, odd, strong, weak and all solar cycles.

is comparable between the strong (~ 37 %) and weak (~ 40 %) cycles for the ap index (Table 4).

Variation of the AE index (Fig. 9) is significantly different than the variations of the Dst and ap indices. In a solar cycle, AE peaks around the descending phase (Fig. 9c). On the yearly basis, the average AE values are enhanced from March/April to September/October. The summer solstice values are significantly higher compared to the winter solstice values. This indicates an annual variation, in agreement with the Lomb–Scargle periodogram analysis result (Fig. 3h). There is no semi-annual variation. The average values during the strong and odd solar cycles are higher compared to the weak and even solar cycles, respectively (Fig. 9a). SC24 exhibited the lowest values of AE compared to other solar cycles (Fig. 9b).

3.2.4 Solar-wind–magnetosphere coupling

The periodogram analysis (Fig. 3j and Table 3) identified a weak annual component in the variations of the solar wind speed V_{sw} (compared with its stronger amplitude longer-




Figure 8. Geomagnetic ap index variation from 1963 to 2019. The panels are in the same format as in Fig. 7.

scale variations). The monthly mean values of V_{sw} during each year of observation are shown in Fig. 10c. In a solar cycle, V_{sw} peaks around the descending phase, indicating a higher occurrence rate of HSSs during this phase. This is also confirmed by the variations of D_{500} (not shown). Interestingly, during the descending phase of SC20, the V_{sw} peak can be noted around March–April; during the SC21 descending phase, two equinoctial peaks are almost symmetric; during the SC22 descending phase, peaks are recorded during the first half of the year; the peaks shift to the second half of the year during the SC23 descending phase; and during the SC24 descending phase, no prominent feature can be inferred. Thus, overall, a shift of the seasonal peak of V_{sw} from the first half to the second half of the year can be observed between the even and the odd cycles. In addition, during the first half of the year, the average values are significantly high during the odd and strong cycles than during the even and weak cycles, respectively (Fig. 10a).

Figure 11 shows the monthly mean values of the coupling function VB_s during all years of observation. In a solar cycle, VB_s peaks around the solar maximum, when almost symmetrical peaks can be observed during the equinoxes and minima during the solstices (Fig. 11c). The lowest values of VB_s are recorded during SC24 (Fig. 11b). There is no prominent difference between the weak and strong cycles, or between the even and odd cycles, except that the February and Octo-




Figure 9. Geomagnetic AE index variation from 1963 to 2019. The panels are in the same format as in Fig. 7.

ber values are higher during the odd and strong cycles compared to those during the even and weak cycles, respectively (Fig. 11a).

4 Conclusions

We used an up-to-date database of substorms, HILDCAAs and geomagnetic storms of varying intensity along with all available geomagnetic indices during the space exploration era (i.e., after 1957) to explore the seasonal features of the geomagnetic activity and their drivers. No such study involving such a long database and all types of geomagnetic activity has been reported before. As substorms, HILDCAAs and magnetic storms of varying intensity have varying solar/interplanetary drivers; such a study is important for a complete understanding of the seasonal features of the geomagnetic response to the solar/interplanetary events. The main findings of this work are discussed below.

First, the semi-annual variation is not a “universal” feature of the geomagnetic activity. While the monthly numbers of substorms and moderate and intense magnetic storms exhibit the semi-annual variation with two equinoctial maxima and a summer solstice minimum, super storms (with a very low occurrence rate) and HILDCAA events do not exhibit any clear seasonal dependence. For geomagnetic in-




Figure 10. Solar wind speed V_{sw} variation from 1963 to 2019. The panels are in the same format as in Fig. 7.




Figure 11. Solar wind coupling function VB_s variation from 1963 to 2019. The panels are in the same format as in Fig. 7.

dices, the monthly mean ring current index Dst and the global geomagnetic activity index ap exhibit the semi-annual variation, while the auroral ionospheric electrojet current index AE exhibits an annual variation with a summer solstice maximum and a winter minimum. These results clearly demonstrate varying solar, interplanetary, magnetospheric and ionospheric processes behind different geomagnetic events and indices. While the magnetic reconnection (Dungey, 1961) between the southward IMF and the northward (dayside) geomagnetic field is the key for any geomagnetic effect, variations in the reconnection process and modulation by other processes may result in different geomagnetic effects (e.g., Gonzalez et al., 1994; Tsurutani et al., 2020; Hajra, 2021a; Hajra et al., 2021, and references therein). In general, major magnetic storms are associated with strong magnetic reconnection continuing for a few hours, while weaker reconnection for an hour or less can cause substorms. On the other hand, discrete and intermittent magnetic reconnection continuing for a long interval of time may lead to HILDCAAs (see Gonzalez et al., 1994, for a detailed comparison).

We observe a clear semi-annual component in the coupling function VB_s , which represents the reconnection electric field or the magnetic flux transfer rate into the magnetosphere. On the other hand, the solar wind speed V_{sw} does not have any semi-annual component, only annual and longer-scale components. As the main focus of the present work is

the seasonal features, we refer the reader to previous works for a discussion on the longer-scale variations in V_{sw} (e.g., Valdés-Galicia et al., 1996; El-Borie, 2002; El-Borie et al., 2020; Hajra, 2021a, c; Hajra et al., 2021, and references therein). However, this result is very interesting. This clearly implies that the solar wind does not have any intrinsic semi-annual variation and that the semi-annual variation in VB_s is due to magnetic configuration (B_s) as suggested previously (e.g., Cortie, 1912; McIntosh, 1959; Boller and Stolov, 1970; Russell and McPherron, 1973). The VB_s semi-annual variation is suggested to cause the semi-annual variations of the substorms, the moderate and intense storms, and the geomagnetic Dst and ap indices. On the other hand, absence of any clear seasonal features in the super storms and HILDCAAs indicates more complex solar wind and magnetic coupling process during these events, which needs further study. As previously established, HILDCAAs are associated with HSSs emanated from the solar coronal holes (e.g., Tsurutani and Gonzalez, 1987; Hajra et al., 2013). Dominating longer-scale variations in V_{sw} (as revealed in the present work) may be a plausible reason for the ~ 4.1 -year variation and lack of any seasonal feature in HILDCAAs (Hajra et al., 2014a; Hajra, 2021c). Annual variation in the auroral ionospheric AE index, as mentioned before, may be attributed to a combined effect of the solar wind V_{sw} variation, the asymmetric pole exposition to the solar radiation, and the ionospheric conduc-

tivity variations (see, e.g., Wang and Lühr, 2007; Tanskanen et al., 2011).

In addition to the above, we found a complex solar activity dependence of the abovementioned seasonal features. The spring–fall asymmetry in substorms and the average V_{sw} variation between the odd and even solar cycles are consistent with results reported by Mursula et al. (2011). An interesting and puzzling result is observed in terms of variations in the semi-annual variability (seasonal modulation between the equinoctial maximum and the solstice minimum) between the strong (odd) and weak (even) solar cycles. While the seasonal modulation in substorms, all storms, moderate storms and the ap index is larger during the weak (and even) solar cycles compared to the strong (and odd) solar cycles, the reverse is true for the intense storms, the Dst index and the coupling function VB_s . At present we do not know the exact mechanism behind this result. In fact, further study is required for a better understanding of the solar cycle dependencies of the geomagnetic activity seasonal features. In conclusion, this study, along with several previous works (e.g., Mursula et al., 2011; Hajra et al., 2013, 2016; Hajra, 2021b), calls for a careful reanalysis of the solar, interplanetary, magnetospheric and ionospheric observations before applying the theoretical semi-annual models.

Data availability. The solar wind plasma and IMF data used in this work are obtained from the OMNI website (<https://omniweb.gsfc.nasa.gov/>, NASA, 2011). The geomagnetic indices are obtained from the World Data Center for Geomagnetism, Kyoto, Japan (<http://wdc.kugi.kyoto-u.ac.jp/>, WDC, 2021). The list of substorms is collected from the SuperMAG website (<https://supermag.jhuapl.edu/>, SuperMAG, 2021). The $F_{10.7}$ solar fluxes are obtained from the Laboratory for Atmospheric and Space Physics (LASP) Interactive Solar Irradiance Data Center (<https://lasp.colorado.edu/lisird/>, LISIRD, 2021).

Author contributions. RH had the original idea. AMSF and RH did the data analysis. RH prepared the first draft. All authors participated in the development, revision of the article and approved the final draft.

Competing interests. The contact author has declared that neither they nor their co-authors have any competing interests.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Acknowledgements. The work of Adriane Marques de Souza Franco is funded by the Brazilian Conselho Nacional de Desenvolvimento Científico e Tecnológico agency

(CNPq, project nos. PQ-300969/2020-1, PQ-301542/2021-0 and PQ-301969/2021-3). The work of Rajkumar Hajra is funded by the Science and Engineering Research Board (SERB, grant no. SB/S2/RJN-080/2018), a statutory body of the Department of Science and Technology (DST), government of India, through a Ramanujan fellowship. Ezequiel Echer would like to thank Brazilian agencies for research grants: CNPq (contract nos. PQ-302583/2015-7 and PQ-301883/2019-0) and Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP, 2018/21657-1). The work of Mauricio José Alves Bolzan was supported by CNPq agency (contract nos. PQ-302330/2015-1, PQ-305692/2018-6) and Fundação de Amparo à Pesquisa do Estado de Goiás agency (FAPEG, contract no. 2012.1026.7000905). We would like to thank two reviewers for extremely valuable suggestions that substantially improved the article.

Financial support. This research has been supported by the Conselho Nacional de Desenvolvimento Científico e Tecnológico (grant nos. PQ-300969/2020-1, PQ-301542/2021-0, PQ-301969/2021-3, PQ-302583/2015-7, PQ-301883/2019-0, PQ-302330/2015-1 and PQ-305692/2018-6), the Science and Engineering Research Board (grant no. SB/S2/RJN-080/2018), the Fundação de Amparo à Pesquisa do Estado de São Paulo (grant no. 2018/21657-1) and the Fundação de Amparo à Pesquisa do Estado de Goiás (grant no. 2012.1026.7000905).

Review statement. This paper was edited by Elias Roussos and reviewed by two anonymous referees.

References

- Akasofu, S.-I.: The development of the auroral substorm, *Planet. Space Sci.*, 12, 273–282, [https://doi.org/10.1016/0032-0633\(64\)90151-5](https://doi.org/10.1016/0032-0633(64)90151-5), 1964.
- Akasofu, S.-I.: Auroral Substorms: Search for Processes Causing the Expansion Phase in Terms of the Electric Current Approach, *Space Sci. Rev.*, 212, 341–381, <https://doi.org/10.1007/s11214-017-0363-7>, 2017.
- Axford, W. I. and Hines, C. O.: A unifying theory of high-latitude geophysical phenomena and geomagnetic storms, *Can. J. Phys.*, 39, 1433–1464, <https://doi.org/10.1139/p61-172>, 1961.
- Baker, D. N., Kanekal, S. G., Pulkkinen, T. I., and Blake, J. B.: Equinoctial and solstitial averages of magnetospheric relativistic electrons: A strong semiannual modulation, *Geophys. Res. Lett.*, 26, 3193–3196, <https://doi.org/10.1029/1999GL003638>, 1999.
- Bartels, J.: Terrestrial-magnetic activity and its relations to solar phenomena, *Terrestrial Magnetism and Atmospheric Electricity*, 37, 1–52, <https://doi.org/10.1029/TE037i001p00001>, 1932.
- Bartels, J.: Twenty-seven day recurrences in terrestrial-magnetic and solar activity, 1923–1933, *Terrestrial Magnetism and Atmospheric Electricity*, 39, 201–202a, <https://doi.org/10.1029/TE039i003p00201>, 1934.
- Boller, B. R. and Stolov, H. L.: Kelvin-Helmholtz instability and the semiannual variation of geomagnetic activity, *J. Geophys. Res.*, 75, 6073–6084, <https://doi.org/10.1029/JA075i031p06073>, 1970.

- Broun, J. A.: Observations in magnetism and meteorology made at Makerstoun in Scotland, *T. Roy. Soc. Edin.-Earth*, 18, 401–402, 1848.
- Burton, R. K., McPherron, R. L., and Russell, C. T.: An empirical relationship between interplanetary conditions and Dst, *J. Geophys. Res.*, 80, 4204–4214, <https://doi.org/10.1029/JA080i031p04204>, 1975.
- Chapman, S. and Ferraro, V. C. A.: A new theory of magnetic storms, *Terrestrial Magnetism and Atmospheric Electricity*, 36, 77–97, <https://doi.org/10.1029/TE036i002p00077>, 1931.
- Cliver, E. W., Kamide, Y., and Ling, A. G.: Mountains versus valleys: Semiannual variation of geomagnetic activity, *J. Geophys. Res.-Space*, 105, 2413–2424, <https://doi.org/10.1029/1999JA900439>, 2000.
- Cliver, E. W., Svalgaard, L., and Ling, A. G.: Origins of the semiannual variation of geomagnetic activity in 1954 and 1996, *Ann. Geophys.*, 22, 93–100, <https://doi.org/10.5194/angeo-22-93-2004>, 2004.
- Cnossen, I. and Richmond, A. D.: How changes in the tilt angle of the geomagnetic dipole affect the coupled magnetosphere–ionosphere–thermosphere system, *J. Geophys. Res.-Space*, 117, A10317, <https://doi.org/10.1029/2012JA018056>, 2012.
- Cortie, A. L.: Sun-spots and Terrestrial Magnetic Phenomena, 1898–1911: the Cause of the Annual Variation in Magnetic Disturbances, *Mon. Not. R. Astron. Soc.*, 73, 52–60, <https://doi.org/10.1093/mnras/73.1.52>, 1912.
- Danilov, A. A., Krymskii, G. F., and Makarov, G. A.: Geomagnetic activity as a reflection of processes in the magnetospheric tail: 1. The source of diurnal and semiannual variations in geomagnetic activity, *Geomagn. Aeron+*, 53, 469–475, <https://doi.org/10.1134/S0016793213040051>, 2013.
- Davis, T. N. and Sugiura, M.: Auroral electrojet activity index AE and its universal time variations, *J. Geophys. Res.*, 71, 785–801, <https://doi.org/10.1029/JZ071i003p00785>, 1966.
- Dungey, J. W.: Interplanetary Magnetic Field and the Auroral Zones, *Phys. Rev. Lett.*, 6, 47–48, <https://doi.org/10.1103/PhysRevLett.6.47>, 1961.
- Durney, B. R.: On the Differences Between Odd and Even Solar Cycles, *Sol. Phys.*, 196, 421–426, <https://doi.org/10.1023/A:1005285315323>, 2000.
- Echer, E., Gonzalez, W. D., Tsurutani, B. T., and Gonzalez, A. L. C.: Interplanetary conditions causing intense geomagnetic storms ($Dst < -100$ nT) during solar cycle 23 (1996–2006), *J. Geophys. Res.*, 113, 1–20, <https://doi.org/10.1029/2007JA012744>, 2008.
- Echer, E., Gonzalez, W. D., and Tsurutani, B. T.: Statistical studies of geomagnetic storms with peak $Dst \leq -50$ nT from 1957 to 2008, *J. Atmos. Sol.-Terr. Phys.*, 73, 1454–1459, <https://doi.org/10.1016/j.jastp.2011.04.021>, 2011.
- El-Borie, M.: On Long-Term Periodicities In The Solar-Wind Ion Density and Speed Measurements During The Period 1973–2000, *Sol. Phys.*, 208, 345–358, <https://doi.org/10.1023/A:1020585822820>, 2002.
- El-Borie, M., El-Taher, A., Thabet, A., and Bishara, A.: The Interconnection between the Periodicities of Solar Wind Parameters Based on the Interplanetary Magnetic Field Polarity (1967–2018): A Cross Wavelet Analysis, *Sol. Phys.*, 295, 122, <https://doi.org/10.1007/s11207-020-01692-2>, 2020.
- Finch, I. D., Lockwood, M. L., and Rouillard, A. P.: Effects of solar wind magnetosphere coupling recorded at different geomagnetic latitudes: Separation of directly-driven and storage/release systems, *Geophys. Res. Lett.*, 35, L21105, <https://doi.org/10.1029/2008GL035399>, 2008.
- Gjerloev, J. W.: The SuperMAG data processing technique, *J. Geophys. Res.*, 117, A09213, <https://doi.org/10.1029/2012JA017683>, 2012.
- Gnevyshev, M. N. and Ohl, A. I.: On the 22-year cycle of solar activity, *Astron. Zh+*, 25, 18–20, 1948.
- Gonzalez, W. D., Joselyn, J. A., Kamide, Y., Kroehl, H. W., Rostoker, G., Tsurutani, B. T., and Vasyliunas, V. M.: What is a geomagnetic storm?, *J. Geophys. Res.*, 99, 5771–5792, <https://doi.org/10.1029/93JA02867>, 1994.
- Gosling, J. T., Bame, S. J., McComas, D. J., and Phillips, J. L.: Coronal mass ejections and large geomagnetic storms, *Geophys. Res. Lett.*, 17, 901–904, <https://doi.org/10.1029/GL017i007p00901>, 1990.
- Grandin, M., Aikio, A. T., and Kozlovsky, A.: Properties and Geoeffectiveness of Solar Wind High-Speed Streams and Stream Interaction Regions During Solar Cycles 23 and 24, *J. Geophys. Res.-Space*, 124, 3871–3892, <https://doi.org/10.1029/2018JA026396>, 2019.
- Guarnieri, F. L.: A study of the solar and interplanetary origin of long-duration and continuous auroral activity events, PhD thesis, INPE, available at: <http://livros01.livrosgratis.com.br/cp012558.pdf> (last access: 23 May 2021), 2005.
- Guarnieri, F. L.: The nature of auroras during high-intensity long-duration continuous AE activity (HILDCAA) events: 1998 to 2001, 235–243, AGU, <https://doi.org/10.1029/167GM19>, 2006.
- Hajra, R.: September 2017 Space-Weather Events: A Study on Magnetic Reconnection and Geoeffectiveness, *Sol. Phys.*, 296, 50, <https://doi.org/10.1007/s11207-021-01803-7>, 2021a.
- Hajra, R.: Seasonal dependence of the Earth’s radiation belt – new insights, *Ann. Geophys.*, 39, 181–187, <https://doi.org/10.5194/angeo-39-181-2021>, 2021b.
- Hajra, R.: Weakest Solar Cycle of the Space Age: A Study on Solar Wind–Magnetosphere Energy Coupling and Geomagnetic Activity, *Sol. Phys.*, 296, 33, <https://doi.org/10.1007/s11207-021-01774-9>, 2021c.
- Hajra, R., Echer, E., Tsurutani, B. T., and Gonzalez, W. D.: Solar cycle dependence of High-Intensity Long-Duration Continuous AE Activity (HILDCAA) events, relativistic electron predictors?, *J. Geophys. Res.-Space*, 118, 5626–5638, <https://doi.org/10.1002/jgra.50530>, 2013.
- Hajra, R., Echer, E., Tsurutani, B. T., and Gonzalez, W. D.: Superposed epoch analyses of HILDCAAs and their interplanetary drivers: Solar cycle and seasonal dependences, *J. Atmos. Sol.-Terr. Phys.*, 121, 24–31, <https://doi.org/10.1016/j.jastp.2014.09.012>, 2014a.
- Hajra, R., Tsurutani, B. T., Echer, E., and Gonzalez, W. D.: Relativistic electron acceleration during high-intensity, long-duration, continuous AE activity (HILDCAA) events: Solar cycle phase dependences, *Geophys. Res. Lett.*, 41, 1876–1881, <https://doi.org/10.1002/2014GL059383>, 2014b.
- Hajra, R., Tsurutani, B. T., Echer, E., Gonzalez, W. D., Brum, C. G. M., Vieira, L. E. A., and Santolik, O.: Relativistic electron acceleration during HILDCAA events: are precursor CIR magnetic storms important?, *Earth Planet. Space*, 67, 109, <https://doi.org/10.1186/s40623-015-0280-5>, 2015a.

- Hajra, R., Tsurutani, B. T., Echer, E., Gonzalez, W. D., and Santolik, O.: Relativistic ($E > 0.6$, > 2.0 , and > 4.0 MeV) electron acceleration at geosynchronous orbit during high-intensity, long-duration, continuous AE activity (HILDCAA) events, *Astrophys. J.*, 799, 39, <https://doi.org/10.1088/0004-637x/799/1/39>, 2015b.
- Hajra, R., Tsurutani, B. T., Echer, E., Gonzalez, W. D., and Gjerloev, J. W.: Supersubstorms (SML < -2500 nT): Magnetic storm and solar cycle dependences, *J. Geophys. Res.-Space*, 121, 7805–7816, <https://doi.org/10.1002/2015JA021835>, 2016.
- Hajra, R., Tsurutani, B. T., and Lakhina, G. S.: The Complex Space Weather Events of 2017 September, *Astrophys. J.*, 899, 3, <https://doi.org/10.3847/1538-4357/aba2c5>, 2020.
- Hajra, R., Franco, A. M. S., Echer, E., and Bolzan, M. J. A.: Long-term variations of the geomagnetic activity: a comparison between the strong and weak solar activity cycles and implications for the space climate, *J. Geophys. Res.-Space*, 126, e2020JA028695, <https://doi.org/10.1029/2020JA028695>, 2021.
- Kanekal, S. G., Baker, D. N., and McPherron, R. L.: On the seasonal dependence of relativistic electron fluxes, *Ann. Geophys.*, 28, 1101–1106, <https://doi.org/10.5194/angeo-28-1101-2010>, 2010.
- Katsavrias, C., Hillaris, A., and Preka-Papadema, P.: A wavelet based approach to Solar–Terrestrial Coupling, *Adv. Space Res.*, 57, 2234–2244, <https://doi.org/10.1016/j.asr.2016.03.001>, 2016.
- Katsavrias, C., Papadimitriou, C., Aminalragia-Giamini, S., Daglis, I. A., Sandberg, I., and Jiggins, P.: On the semi-annual variation of relativistic electrons in the outer radiation belt, *Ann. Geophys.*, 39, 413–425, <https://doi.org/10.5194/angeo-39-413-2021>, 2021.
- Lakhina, G. S. and Tsurutani, B. T.: Chapter 7 – Supergeomagnetic Storms: Past, Present, and Future, in: *Extreme Events in Geospace: Origins, Predictability, and Consequences*, edited by: Buzulukova, N., 157–185, Elsevier, Amsterdam, the Netherlands, <https://doi.org/10.1016/B978-0-12-812700-1.00007-8>, 2018.
- Lamy, P. L., Floyd, O., Boclet, B., Wojak, J., Gilardy, H., and Barlyaeva, T.: Coronal Mass Ejections over Solar Cycles 23 and 24, *Space Sci. Rev.*, 215, 39, <https://doi.org/10.1007/s11214-019-0605-y>, 2019.
- Le Mouél, J.-L., Blanter, E., Chulliat, A., and Shnirman, M.: On the semiannual and annual variations of geomagnetic activity and components, *Ann. Geophys.*, 22, 3583–3588, <https://doi.org/10.5194/angeo-22-3583-2004>, 2004.
- Li, X., Baker, D. N., Kanekal, S. G., Looper, M., and Temerin, M.: Long term measurements of radiation belts by SAMPEX and their variations, *Geophys. Res. Lett.*, 28, 3827–3830, <https://doi.org/10.1029/2001GL013586>, 2001.
- LISIRD: LASP Interactive Solar Irradiance Data Center, available at: <https://lasp.colorado.edu/lisird/>, last access: 23 May 2021.
- Lockwood, M., Owens, M. J., Barnard, L. A., Haines, C., Scott, C. J., McWilliams, K. A., and Coxon, J. C.: Semi-annual, annual and Universal Time variations in the magnetosphere and in geomagnetic activity: 1. Geomagnetic data, *J. Space Weather Spac.*, 10, 23, <https://doi.org/10.1051/swsc/2020023>, 2020.
- Lomb, N. R.: Least-squares frequency analysis of unequally spaced data, *Astrophys. Space Sci.*, 39, 447–462, <https://doi.org/10.1007/BF00648343>, 1976.
- Marques de Souza, A., Echer, E., Bolzan, M. J. A., and Hajra, R.: Cross-correlation and cross-wavelet analyses of the solar wind IMF B_z and auroral electrojet index AE coupling during HILDCAAs, *Ann. Geophys.*, 36, 205–211, <https://doi.org/10.5194/angeo-36-205-2018>, 2018.
- McIntosh, D. H.: On the annual variation of magnetic disturbance, *Philos. T. R. Soc. S-A*, 251, 525–552, <https://doi.org/10.1098/rsta.1959.0010>, 1959.
- McPherron, R. L. and Chu, X.: The Midlatitude Positive Bay Index and the Statistics of Substorm Occurrence, *J. Geophys. Res.*, 123, 2831–2850, <https://doi.org/10.1002/2017JA024766>, 2018.
- Mendes, O., Domingues, M. O., Echer, E., Hajra, R., and Menconi, V. E.: Characterization of high-intensity, long-duration continuous auroral activity (HILDCAA) events using recurrence quantification analysis, *Nonlin. Processes Geophys.*, 24, 407–417, <https://doi.org/10.5194/npg-24-407-2017>, 2017.
- Meng, C.-I., Mauk, B., and McIlwain, C.: Electron precipitation of evening diffuse aurora and its conjugate electron fluxes near the magnetospheric equator, *J. Geophys. Res.*, 84, 2545–2558, <https://doi.org/10.1029/JA084iA06p02545>, 1979.
- Mursula, K. and Zieger, B.: Long-term north-south asymmetry in solar wind speed inferred from geomagnetic activity: A new type of century-scale solar oscillation?, *Geophys. Res. Lett.*, 28, 95–98, <https://doi.org/10.1029/2000GL011880>, 2001.
- Mursula, K., Hiltula, T., and Zieger, B.: Latitudinal gradients of solar wind speed around the ecliptic: Systematic displacement of the streamer belt, *Geophys. Res. Lett.*, 29, 1–4, <https://doi.org/10.1029/2002GI015318>, 2002.
- Mursula, K., Tanskanen, E., and Love, J. J.: Spring-fall asymmetry of substorm strength, geomagnetic activity and solar wind: Implications for semiannual variation and solar hemispheric asymmetry, *Geophys. Res. Lett.*, 38, L06104, <https://doi.org/10.1029/2011GL046751>, 2011.
- Nakagawa, Y., Nozawa, S., and Shinbori, A.: Relationship between the low-latitude coronal hole area, solar wind velocity, and geomagnetic activity during solar cycles 23 and 24, *Earth Planet. Space*, 71, 24, <https://doi.org/10.1186/s40623-019-1005-y>, 2019.
- NASA: OMNIWeb Plus, available at: <https://omniweb.gsfc.nasa.gov/>, last access: 23 May 2021.
- Newell, P. T. and Gjerloev, J. W.: Evaluation of SuperMAG auroral electrojet indices as indicators of substorms and auroral power, *J. Geophys. Res.*, 116, A12211, <https://doi.org/10.1029/2011JA016779>, 2011.
- Newton, H. W. and Nunn, M. L.: The Sun's Rotation Derived from Sunspots 1934–1944 and Additional Results, *Mon. Not. R. Astron. Soc.*, 111, 413–421, <https://doi.org/10.1093/mnras/111.4.413>, 1951.
- Nykyri, K., Bengtson, M., Angelopoulos, V., Nishimura, Y., and Wing, S.: Can Enhanced Flux Loading by High-Speed Jets Lead to a Substorm? Multipoint Detection of the Christmas Day Substorm Onset at 08:17 UT, 2015, *J. Geophys. Res.*, 124, 4314–4340, <https://doi.org/10.1029/2018JA026357>, 2019.
- O'Brien, T. P. and McPherron, R. L.: Seasonal and diurnal variation of Dst dynamics, *J. Geophys. Res.-Space*, 107, SMP 3–1–SMP 3–10, <https://doi.org/10.1029/2002JA009435>, 2002.
- Owens, M. J., Lockwood, M., Barnard, L. A., Scott, C. J., Haines, C., and Macneil, A.: Extreme Space-Weather Events and the Solar Cycle, *Sol. Phys.*, 296, 82, <https://doi.org/10.1007/s11207-021-01831-3>, 2021.
- Perreault, P. and Akasofu, S.-I.: A study of geomagnetic storms, *Geophys. J. Roy. Astr. Soc.*, 54, 547–573, <https://doi.org/10.1111/j.1365-246X.1978.tb05494.x>, 1978.

- Rawat, R., Echer, E., and Gonzalez, W. D.: How Different Are the Solar Wind–Interplanetary Conditions and the Consequent Geomagnetic Activity During the Ascending and Early Descending Phases of the Solar Cycles 23 and 24?, *J. Geophys. Res.-Space*, 123, 6621–6638, <https://doi.org/10.1029/2018JA025683>, 2018.
- Richardson, I. G., Cane, H. V., and Cliver, E. W.: Sources of geomagnetic activity during nearly three solar cycles (1972–2000), *J. Geophys. Res.-Space*, 107, SSH 8–1–SSH 8–13, <https://doi.org/10.1029/2001JA000504>, 2002.
- Rostoker, G.: Geomagnetic indices, *Rev. Geophys.*, 10, 935–950, <https://doi.org/10.1029/RG010i004p00935>, 1972.
- Rostoker, G.: Identification of substorm expansive phase onsets, *J. Geophys. Res.*, 107, SMP 26–1–SMP 26–9, <https://doi.org/10.1029/2001JA003504>, 2002.
- Russell, C. T.: Geophysical Coordinate Transformations, chap. Appendix 3, 184–196, Cambridge University Press, Cambridge, UK, <https://doi.org/10.1017/9781139878296.019>, 1971.
- Russell, C. T. and McPherron, R. L.: Semiannual variation of geomagnetic activity, *J. Geophys. Res.*, 78, 92–108, <https://doi.org/10.1029/JA078i001p00092>, 1973.
- Sabine, E.: VIII. On periodical laws discoverable in the mean effects of the larger magnetic disturbance. 2014; No. II, *Philos. T. R. Soc. Lond.*, 142, 103–124, <https://doi.org/10.1098/rstl.1852.0009>, 1852.
- Scargle, J. D.: Studies in astronomical time series analysis. II. Statistical aspects of spectral analysis of unevenly spaced data, *Astrophys. J.*, 263, 835–853, 1982.
- Schwabe, H.: Sonnen-Beobachtungen im Jahre 1843, *Astron. Nachr.*, 21, 233–236, 1844.
- Scolini, C., Messerotti, M., Poedts, S., and Rodriguez, L.: Halo coronal mass ejections during Solar Cycle 24: reconstruction of the global scenario and geoeffectiveness, *J. Space Weather Spac.*, 8, A09, <https://doi.org/10.1051/swsc/2017046>, 2018.
- Souza, A. M., Echer, E., Bolzan, M. J. A., and Hajra, R.: A study on the main periodicities in interplanetary magnetic field Bz component and geomagnetic AE index during HILDCAA events using wavelet analysis, *J. Atmos. Sol.-Terr. Phys.*, 149, 81–86, <https://doi.org/10.1016/j.jastp.2016.09.006>, 2016.
- Sugiura, M.: Hourly values of equatorial Dst for the IGY, *Ann. Intern. Geophys. Year*, 35, 9, 1964.
- SuperMAG: <https://supermag.jhuapl.edu/>, last access: 23 May 2021.
- Syed Ibrahim, M., Joshi, B., Cho, K.-S., Kim, R.-S., and Moon, Y.-J.: Interplanetary Coronal Mass Ejections During Solar Cycles 23 and 24: Sun–Earth Propagation Characteristics and Consequences at the Near-Earth Region, *Sol. Phys.*, 294, 54, <https://doi.org/10.1007/s11207-019-1443-5>, 2019.
- Tanskanen, E. I., Pulkkinen, T. I., Viljanen, A., Mursula, K., Partamies, N., and Slavin, J. A.: From space weather toward space climate time scales: Substorm analysis from 1993 to 2008, *J. Geophys. Res.*, 116, A00134, <https://doi.org/10.1029/2010JA015788>, 2011.
- Tapping, K. F.: Recent solar radio astronomy at centimeter wavelengths: The temporal variability of the 10.7-cm flux, *J. Geophys. Res. Atmos.*, 92, 829–838, <https://doi.org/10.1029/JD092iD01p00829>, 1987.
- Thorne, R. M., Ni, B., Tao, X., Horne, R. B., and Meredith, N. P.: Scattering by chorus waves as the dominant cause of diffuse auroral precipitation, *Nature*, 467, 943–946, <https://doi.org/10.1038/nature09467>, 2010.
- Tsurutani, B. T. and Gonzalez, W. D.: The cause of high-intensity long-duration continuous AE activity (HILDCAAs): Interplanetary Alfvén wave trains, *Planet. Space Sci.*, 35, 405–412, [https://doi.org/10.1016/0032-0633\(87\)90097-3](https://doi.org/10.1016/0032-0633(87)90097-3), 1987.
- Tsurutani, B. T. and Meng, C.-I.: Interplanetary magnetic-field variations and substorm activity, *J. Geophys. Res.*, 77, 2964–2970, <https://doi.org/10.1029/JA077i016p02964>, 1972.
- Tsurutani, B. T., Gonzalez, W. D., Tang, F., Akasofu, S. I., and Smith, E. J.: Origin of interplanetary southward magnetic fields responsible for major magnetic storms near solar maximum (1978–1979), *J. Geophys. Res.-Space*, 93, 8519–8531, <https://doi.org/10.1029/JA093iA08p08519>, 1988.
- Tsurutani, B. T., Gonzalez, W. D., Tang, F., and Lee, Y. T.: Great magnetic storms, *Geophys. Res. Lett.*, 19, 73–76, <https://doi.org/10.1029/91GL02783>, 1992.
- Tsurutani, B. T., Gonzalez, W. D., Guarnieri, F., Kamide, Y., Zhou, X., and Arballo, J. K.: Are high-intensity long-duration continuous AE activity (HILDCAA) events substorm expansion events?, *J. Atmos. Sol.-Terr. Phys.*, 66, 167–176, <https://doi.org/10.1016/j.jastp.2003.08.015>, 2004.
- Tsurutani, B. T., Hajra, R., Echer, E., and Lakhina, G. S.: Comment on “First Observation of Mesosphere Response to the Solar Wind High-Speed Streams” by W. Yi et al., *J. Geophys. Res.*, 124, 8165–8168, <https://doi.org/10.1029/2018JA026447>, 2019.
- Tsurutani, B. T., Lakhina, G. S., and Hajra, R.: The physics of space weather/solar-terrestrial physics (STP): what we know now and what the current and future challenges are, *Nonlin. Processes Geophys.*, 27, 75–119, <https://doi.org/10.5194/npg-27-75-2020>, 2020.
- Valdés-Galicia, J. F., Pérez-Enríquez, R., and Otaola, J. A.: The Cosmic-Ray 1.68-Year Variation: a Clue to Understand the Nature of the Solar Cycle?, *Sol. Phys.*, 167, 409–417, <https://doi.org/10.1007/BF00146349>, 1996.
- von Humboldt, A.: Die vollständigste aller bisherigen Beobachtungen über den Einfluss des Nordlichts auf die Magnetnadel angestellt, *Ann. Phys.*, 29, 425–429, <https://doi.org/10.1002/andp.18080290806>, 1808.
- Waldmeier, M.: Neue Eigenschaften der Sonnenfleckenkurve, *Astronomische Mitteilungen der Eidgenössischen Sternwarte Zürich*, 14, 105–136, 1934.
- Wang, H. and Lüher, H.: Seasonal-longitudinal variation of substorm occurrence frequency: Evidence for ionospheric control, *Geophys. Res. Lett.*, 34, L07 104, <https://doi.org/10.1029/2007GL029423>, 2007.
- WDC: World Data Center for Geomagnetism, Kyoto, available at: <http://wdc.kugi.kyoto-u.ac.jp/>, last access: 23 May 2021.
- Wilson, R. M.: Bimodality and the Hale cycle, *Sol. Phys.*, 117, 269–278, <https://doi.org/10.1007/BF00147248>, 1988.